USAARL REPORT NO. 75-9

EVALUATION OF PROPOSED ELECTROPLATED HGU-4/P FRAMES

Ву

MAJ Roger W. Wiley, O.D., Ph.D. LTC Frank S. Pettyjohn, M.D. MAJ David D. Glick, O.D.

US Army Aeromedical Research Laboratory Fort Rucker, Alabama 36360

February 1975

Final Report

This document has been approved for public release and sale, its distribution is unlimited.



SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM		
1 REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER		
75 9				
4. TITLE (and Subtitle) EVALUATION OF PROPOSED ELECTROPLAT	mp um A/D	5. TYPE OF REPORT & PERIOD COVERED		
FRAMES	ED MGU-4/E	Final		
LIAA-0	ļ	6. PERFORMING DRG. REPORT NUMBER		
	<u> </u>			
7. AUTHOR(a)		8. CONTRACT OR GRANT NUMBER(a)		
MAJ Roger W. Wiley, O.D., Ph.D.	ļ			
LTC Frank S. Pettyjohn, M.D.	!			
MAJ David D. Glick, O.D. P. Performing organization name and address		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT HUMBERS		
Bio-Optics Division (SGRD-UAC)	!			
U S Army Aeromedical Research Labo	ratory	DD Form 1498		
Fort Rucker, AL 36360		12. REPORT DAYE		
U S Army Aeromedical Research Labo	ratory			
SGRD-UAC		February 1975 13. NUMBER OF PAGES		
Fort Rucker, AL 36360 14. MONITORING AGENCY NAME & ADDRESS(II differen	O Seller Office)	9 15. SECURITY CLASS. (of this report)		
		15. SECURITY CLASS, (or and report)		
USA Medical Research and Developme Washington, DC 20314	nt Command	Unclassified		
wasittiigcoit, DC 20014		IS. DECLASSIFICATION/DOWNGRADING		
16. DISTRIBUTION STATEMENT (of this Report)		3011 2002 2		
This document has been approved for public release and sale; its distribution is unlimited.				
17. DISTRIBUTION STATEMENT (at the abstract entered	to Disch 20 II different fre	- Sancet		
17. DISTRIBUTION STATEMENT (or the economic con	IN 8106R 49, 11 6	яп Керопу		
		i		
14. SUPPLEMENTARY NOTES				
18. SUPPLEMENTANT NOTES		!		
		1		
		· ·		
19. KEY WORDS (Continue on reverse side if necessary and	nd Identify by block number;)		
Nickel dermatitis		•		
Skin reaction				
Contact dermatitis Aviator frames				
Aviator frames Nickel-containing frames		ı		
20. ABSTRACT (Continue on reverse side if necessary and	d identify by block number)			
A gold electroplated frame has	been recommended	ed to replace the standard		
gold-filled aviator frame. Since	the proposed fram	me contains a nickel-silver		
based metal, the frame was evaluated under field and laboratory conditions at the U.S. Army Aeromedical Research Laboratory. Of the 18 subjects who				
wore the test frames for three mon	ths, one subject	, an aviator, developed		
a mild dermatitis along the fronta	d and supraorbita	al portion of the face.		
Chemical analysis indicated "free" nickel in sufficient quantity to cause a				

EDITION OF 1 NOV 65 IS OBSOLETE

DD 1 JAN 75 1473

Unclassified

UNCIASSIFIED SECURITY CLASSIFICATION OF THIS PAGE(When Date Entered)
reaction from nickel sensitive individuals. This study has shown that some skin reaction can be expected from a small percentage of wearers if the gold electroplated frame replaces the gold-filled frame.

NOTICE

Qualified requesters may obtain copies from the Defense Documentation Center (DDC), Cameron Station, Alexandria, Virginia. Orders will be expedited if placed through the librarian or other person designated to request documents from DDC (Formerly ASTIA).

Change of Address

Organization receiving reports from the US Army Aeromedical Research Laboratory on automatic mailing lists should confirm correct address when corresponding about laboratory reports.

Disposition

Destroy this report when it is no longer needed. Do not return it to the originator.

Disclaimer

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

SUMMARY

A gold electroplated frame has been recommended to replace the standard gold-filled aviator frame. Since the proposed frame contains a nickel-silver based metal, the frame was evaluated under field and laboratory conditions at the U.S. Army Aeromedical Research Laboratory. Of the 18 subjects who wore the test frames for three months, one subject, an aviator, developed a mild dermatitis along the frontal and supraorbital portion of the face. Chemical analysis indicated "free" nickel in sufficient quantity to cause a reaction from nickel sensitive individuals. This study has shown that some skin reaction can be expected from a small percentage of wearers if the gold electroplated frame replaces the gold-filled frame.

Approved:

ROBERT W. BAI Colonel, MSC

Commanding

TABLE OF CONTENTS

	PAGE
SUNTIARY	
INTRODUCTION	1
METHODS	2
RESULTS	3
Clinical Evaluation	3
Frame Evaluation	4
DISCUSSION	4
RECOMMENDATIONS	5
REFERENCES	7
APPENDIX - Questionnaire	A-1

INTRODUCTION

Because of the considerable increase in the price of gold during the past several years, the manufacturers supplying the aviator gold-filled ophthalmic frame recently notified the Defense Personnel Support Center (DPSC) that fabrication of this frame was no longer cost-effective, and they were, therefore, unable to support the Army's requirements for these frames. As an alternative metal frame, the civilian manufacturers have recommended that the DPSC purchase an ophthalmic frame consisting of a nickel-silver base metal, plated with bright nickel, and overplated with gold. However, the Army dispensed nickel-containing frames in the 1940's and 1950's with the resulting experience of many soldiers reporting a dermatitis around the area of contact of the frames with the skin.

Nickel has been a well recognized source of metal contact dermatitis. 1,2,3 The clinical picture is an eczematous dermatitis with pruritus, scaling and erythema progressing to lichenification. Common offending items containing nickel include rings, earrings, jewelry, spectacle frames, wrist watches, suspenders, and support garments.

Nickel sensitivity is considered more common in women than men. In the United Kingdom and Scandanavia, nickel is the most common cause of contact dermatitis in women. Women, by their use of jewelry, do provide for greater exposure of nickel to skin contact than men. In men, nickel dermatitis occurs most commonly due to occupational exposure. In 60% of the cases, the eruption begins on the hands. Sensitization by wrist watches and spectacles is common.

Chrome plating has been used to decrease available surface nickel. Dermatitis has been produced by nickel "leaking out" through pores of the chromium plating. Sweating has been implicated in the production of dermatitis as the sodium chloride may combine with nickel-silver to give silver chloride. This process provides the basis for the finding of increasingly severe dermatitis in persons who perspire profusely.

The problem of nickel sensitivity should be considered in any patient with a metallic device or prosthesis externally or internally applied. Implanted stainless steel screws having 14% nickel have produced a severe dermatitis. A Wainwright stainless steel orthopedic plate containing 8 to 12% nickel and 17 to 20% chromium produced a dermatitis and "rejection" of the implant.

Interestingly, a known nickel sensitive patient has received a Bjork-Shiley mitral valve prosthesis containing three grams of metal of which 10% is nickel without reaction after two years. It is

suggested that the small surface area, and thus less metal, may prevent difficulty due to constant wear; however, nickel may be delivered over a prolonged period. 5

The use of nickel-gold alloys and nickel plated earrings following ear piercing has been documented as a cause of contact dermatitis.

The presence of nickel in contact with the skin in amounts of 10 to 14% would indicate a strong possibility of allergic reaction in a sensitive individual. The plating specification for the U.S. Government HGU-4/P flying goggles provide a base metal of not less than 14% nickel silver, with nickel electroplate of not less than 50 millionths of an inch in thickness and overplated with a minimum of 60 millionths of an inch of 23K electrodeposit gold. Because of the possibility of contact dermatitis despite the buffer layer provided by the gold electrodeposit, the proposed frames were evaluated under field and laboratory conditions at the U.S. Army Aeromedical Research laboratory.

METHODS

The sample frames used in this study were supplied through the Defense Personnel Support Center at the request of the American Optical Corporation. Of the twenty frames received from DPSC, eighteen pairs were used in the field study, one pair was retained for laboratory chemical analysis, and one pair was forwarded to the Medical Optical Laboratory, Fitzsimons Army Medical Center.

The subjects who volunteered to wear the test frames for the field study were selected as being reasonably representative of the potential user population. Table 1 shows the number of subjects within each of

	PRESCRIPTION LENSES	PLANO LENSES	
AVIATOR	4	÷	7
TECHNICAL OBSERVER FLIGHT STATUS	5	2	7
NOT ON FLIGHT STATUS		1	4
	9	0	

Table 1. The number of subjects within each of the test groups.

the various groups. Included in the non-prescription, non-flight status group was one female and one black subject; all other subjects were white males. Before dispensing the test spectacles, each subject was individually briefed concerning the purposes of the study, the possible skin reactions to the frame, and what observations he or she should make periodically while wearing the test frame. Depending upon each subject's visual requirements, prescription lenses or plano tinted lenses were mounted into the frames. Each frame was adjusted to allow normal fitting characteristics, and the subjects were instructed to wear the test frames as they would normally wear their glasses.

After the subjects had worn the frames for approximately three months, they returned to the Laboratory to complete a questionnaire (appended). Each subject was examined by a flight surgeon as required, and the frames were examined by an optometrist for wear and durability.

Several of the frames were tested for the presence of "free" nickel with a 1% alcohol solution of dimethylglyoxime (DMG) with ammonia (furnished by Westwood Pharmaceuticals, Inc., Buffalo, New York). This precipitate test will detect the presence of "free" nickel in sufficient concentration (1:10,000) to produce dermatitis in nickelsensitive individuals.

RESULTS

Clinical Evaluation

In our series of 18 subjects wearing the proposed HGU-4/P flying goggles, one subject, an aviator, developed a mild eczematous eruption of the frontal and supraorbital portion of the face. The area was erythematous with mild scaling. Pruritus was not described. The onset of the crythema was within two hours of wearing the frames and, reportedly, occurred each time the frame was worn while performing flying duties.

This aviator was emmetropic and usually wore his sunglasses only while flying. At the time of initial report of a dermatitis reaction, he had worn the standard issue gold-filled flying goggles for approximately seven years without experiencing any dermatitis from the frames. Subsequent to the test period, a trial of wearing the standard gold-filled frames with identical wearing conditions produced no reaction. Following this period, a second trial of prolonged (eight hours) wear of the test electroplated frames also failed to produce any reaction. However, several variations from the original wearing conditions should be emphasized. For this second trial, the sunglass lenses were removed

to facilitate indoor wear and the frames were considerably lighter which would affect the bearing areas on the face. Also, and probably of most significance, this second trial was conducted in an office area so that true cockpit conditions, including wear of the SPH-4 helmet with consequent perspiration, were not approximated. The possiblity that the presence of perspiration contributed to the reaction was increased in that skin testing of this patient utilizing a "buffalo" nickel coin failed to present any reaction after 24 hours contact.

Frame Evaluation

Four frames were analyzed with the dimethylglyoxime spot test for the presence of "free" nickel. One of these frames had never been worn while the remaining three, including the pair resulting in dermatitis, had been worn for approximately three months. The frames were analyzed along the bridge, eyewire, and temple. All of the frames tested positive for the presence of "free" nickel in sufficient quantity to cause an allergic reaction from nickel-sensitive individuals. In addition, one standard gold-filled frame which had been worn previously was tested, and the results were negative for "free" nickel.

In general, the condition of the frames were as might be expected after three months wear. However, a significant percentage (39%) of them showed pitting and etching of the metal core underlying the clear plastic temple paddles. This might indicate a defect in the manufacturing process in which the seal around the plastic paddle temple cover was inadequate. Perspiration had probably been allowed to seep underneath the paddle and had pitted the metal core.

DISCUSSION

The results of this study provide evidence of "free" nickel arising from a nickel base of a known sensitizing concentration of not less than 14% as specified for the proposed HGU-4/P flying goggles. Chemical analysis has shown the potential for reaction, and field testing has developed one borderline positive dermatitis. The series of this study, although small, would indicate a 5% -6% incidence of nickel reaction.

The one dermatitis reaction was interesting in that the subject failed to develop similar or more pronounced symptoms with prolonged wear, and did not react to the "buffalo" nickel. However, as discussed earlier, the variations in the wearing conditions might explain the failure to develop symptoms with the longer wearing schedule. With the lenses removed, the frames were much lighter. Also, the frames

were worn only in an air-conditioned office. The hot and humid cockpit environment quite possibly causes sufficient perspiration to interact with the metal frame to result in the skin reaction.

The necessary restrictions imposed on this study must be recognized in considering the significance of these results. The sample size (n = 18) was quite small, and the test period only extended for three months. However, these restrictions, if anything, force a conservative estimate of the magnitude of the dermatitis problems to be expected if these frames are dispensed to military personnel. No attempt was made to screen the subjects for prior history of any dermatological conditions. When the study was terminated, three subjects reported that they had experienced a metal dermatitis previously, and none of these three evidenced any reaction to the test frames. However, these three subjects had only worn the frames in air-conditioned environments.

A problem that was not realistically addressed with this evaluation is the question of the durability of the gold electroplating. Since this study had to be completed within a relatively short time period, the frames were inspected after only three months use. Even after this short period, there was some evidence of rubbing along the inside of the temples and across the top of the eyewires with some of the frames. If a sufficient amount of the gold electroplating finish rubbed off to expose the nickel base, the dermatitis reactions might have been more frequently experienced.

RECOMMENDATIONS

- 1. Additional effort should be made to find a frame to replace the HCU-4/P flying goggles. Although the problems experienced during this evaluation were relatively minor, these results indicate that some reactions probably will be experienced by nickel-sensitive individuals. This is particularly so for those individuals working in warm environments, where perspiration might interact with the nickel-silver to increase the possibility and/or severity of reaction.
- 2. If an acceptable alternative frame cannot be found for the standard gold-filled frame, appropriate medical personnel (e.g. flight surgeons, eye clinics, dermatology clinics, etc.) should receive information concerning the potential skin reactions. In addition, a warning notice should be included with each pair of the gold electroplated frames issued to alert user personnel of the potential problems.
- 3. A sufficient supply of the standard gold-filled frames should be maintained for those personnel who might suffer from nickel dermatitis.

4. The U.S. Army Aeromedical Research Laboratory would encourage formation of a tri-service working group to develop a mutually acceptable alternative frame to replace the metal HGU-4/P flying goggles.

REFERENCES

- 1. Rook, A., Wilkinson, D.S., Ebling, F.J.G.: Textbook of Dermatology, 2nd Ed., Vol. 1. Blackwell Scientific Publication, London, 1972.
- 2. Domonkos, A.N.: Diseases of the Skin. W.B. Saunders Co., Philadelphia, PA, 1971.
- 3. Fisher, A.A.: <u>Contact Dermatitis</u>. Lea Febiger, Philadephia, 1967.
- 4. Barranco, V.P., Soloman, H.: "Eczematous Dermatitis from Nickel". JAMA, 220: 1244, 1972.
- 5. Pegum, J.S.: "Nickel Allergy". Lancet, Vol. 1, 674, 1974.
- 6. Lyell, A., Bain, W.H.: "Nickel Allergy and Valve Replacement".
 Lancet, Vol. 1, 408-409, 1974.
- 7. Fisher, A.A.: "Ear Piercing Hazard of Nickel-Gold Sensitization".

 JAMA, 228: 1226, 1974.
- 8. Plating Specification/Recommendation for U.S. Government HGU-4/P Flying Goggle.

AVIATION LYECLASS FRAME STUDY

	NAME	RANK	SSN	
	ORGANI ZATION RA	CE:	HAIR COL	NR .
1.	Issued: clear lens sun	glasses		
2.	Do you wear glasses full time?	part	time?	
	How many hours/day?			
3.	Are you allergic to:			
	a. Medication?		Yes	No
	b. Any chemicals?			\Box
	If so, list			
	c. Any soaps or detergents?			$\overline{\Box}$
	If so, list			
	d. Any type clothing or cloth?		\Box	$\overline{\Box}$
	If so, what type?	• • • • • • • • • • • • • • • • • • • •		
4.	Have you ever had a skin rash from	your shoes?		
5.	Have you ever developed a rash fro watch band?	m your wrist		
6.	Have you ever developed a rash fro in contact with your skin?	m any metal		
7.	Since wearing the experimental eye have you noted any rash or itching the face?	glass frames on or aroun	d d	
	a. Your nose?			
	b. Your eyehrows?		$\overline{\Box}$	<u>/</u> 7 <u>/</u> 7

	c. Your hairline?	$\overline{//}$	\Box 7
8.	Are the experimental eyeglass frames in your opinion		
	Comfortable?		\Box
	Provide good fit?	$\overline{\Box}$	\Box
Q	Any comments?		